

# Product Sheet

## H\_FOLH1(PSMA) PC-3 Cell Line

Catalog number: GM-C35058

Version 3.3.1.260323

<b>Description</b>	H_FOLH1(PSMA) PC-3 Cell Line is a clonal stable PC-3 cell line that constitutively expresses the human FOLH1(PSMA) gene, constructed using lentiviral technology.
<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	H_FOLH1(PSMA)
<b>Gene ID/Uniprot ID</b>	Q04609-1 1994-06-01 v1
<b>Host Cell</b>	PC-3 Cell Line
<b>Recovery Medium</b>	F12K+10% FBS+1% P.S
<b>Growth medium</b>	F12K+10% FBS+1% P.S+0.5 µg/mL Puromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
Puromycin	Genomeditech/ <a href="#">GM-040401</a>
Pen/Strep	Thermo/15140-122
Fetal Bovine Serum	ExCell/FSP500
F12K	BOSTER/PYG0036
Anti-FOLH1(PSMA) hIgG1 Reference Antibody (Rosobio)	Genomeditech/ <a href="#">GM-87705MAB</a>

## Figures




Figure 1 | H\_FOLH1(PSMA) PC-3 Cell Line (Cat. GM-C35058) was determined by flow cytometry using Anti-FOLH1(PSMA) hIgG1 Reference Antibody (Rosobio) (Cat. [GM-87705MAB](#)).




Figure 2 | Tumor growth curves of H\_FOLH1(PSMA) PC-3 in BALB/c nude mice. H\_FOLH1(PSMA) PC-3 cells ( $1 \times 10^6$  per mouse) were subcutaneously inoculated into BALB/c nude mice (male, 8 weeks old,  $n = 3$ ). Tumor volume was measured twice per week and is presented as mean  $\pm$  SEM.




Figure 3 | Body weight changes after implantation of H\_FOLH1(PSMA) PC-3 in BALB/c nude mice. Under the same conditions, body weight was measured twice per week and is presented as mean  $\pm$  SEM.

## Cell Recovery

Recovery Medium: F12K+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at  $-70^{\circ}\text{C}$ . Storage at  $-70^{\circ}\text{C}$  will result in loss of viability.

- Thaw the vial by gentle agitation in a  $37^{\circ}\text{C}$  water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately  $176 \times g$  for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- Incubate the culture at  $37^{\circ}\text{C}$  in a suitable incubator. A 5%  $\text{CO}_2$  in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at  $176 \times g$  for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to  $3 \times 10^6$  cells/mL.
- Aliquot 1 mL into each vial.

- d) Place the vial in a controlled-rate freezing container and store at  $-80^{\circ}\text{C}$  for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: F12K+10% FBS+1% P.S+0.5  $\mu\text{g}/\text{mL}$  Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Subculturing is necessary when the cell density reaches 90%. It is recommended to perform subculturing at a ratio of 1:3 to 1:4 every 2-3 days. Ensure that the density does not exceed 90%, as overcrowding can lead to reduced viability due to compression.
- b) Remove and discard culture medium.
- c) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- d) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 3 minutes at  $37^{\circ}\text{C}$ ).
- e) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at  $37^{\circ}\text{C}$  to facilitate dispersal.
- f) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- g) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- h) Incubate cultures at  $37^{\circ}\text{C}$ .

**Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- a) It is normal to observe a number of dead cells immediately after thawing. The condition will improve significantly after adjustment. Once the cells stabilize, the number of dead cells will decrease after subculturing, and the cell growth rate will become stable.
- b) Ensure that the cell density does not exceed 90%, as overcrowding may lead to reduced viability due to compression.
- c) FBS should be heat-inactivated at  $56^{\circ}\text{C}$  for 30 minutes to inactivate complement and certain viruses, with minimal impact on most growth factor and cytokine activities.

## Sequence

PSMA Q04609-1

MWNLLHETDSAVATARRPRWLCAGALVLAGGFFLLGFLFGWFIKSSNEATNITPKHNMKAFDELKAENIK  
KFLYNFTQIPHLAGTEQNFQLAKQIQSQWKEFGLDSVELAHYDVLLSYPNKTHPNYISIINEDGNEIFNTSLFEP  
PPPGYENVSDIVPPFSAFSPQGMPEGDLVYVNYARTEDFFKLERDMKINCSGKIVIARYGKVFRGNKVKNAQ  
LAGAKGVILYSDPADYFAPGVKSYPDGWNLPGGGVQGRNINLNGAGDPLTPGYANEYAYRRGIAEAVGL  
PSIPVHPIGYDAQKLEKMGGSAPPDSSWRGSLKVPYNVGPGFTGNFSTQKVKMHIHSTNEVTRIYNVIGTL

RGAVEPDRYVILGGHRDSWVFGGIDPQSGAAVVHEIVRSFGTLKKEGWRPRRILFASWDAEEFGLLGSTEW  
 AEENSRLQLQERGVAYINADSSIEGNYTLRVDCTPLMYSLVHNLTKELKSPDEGFEGKSLYESWTKKSPSPEFS  
 GMPRISKLGSGNDFEVFFQRLGIASGRARYTKNWETNKFSGYPLYHSVYETYELVEKFYDPMFKYHLTVAQ  
 VRGGMVFELANSIVLPFDCRDYAVVLRKYADKIYSISMKHPQEMKTYSVSFDLSFAVKNFTEIASKFSERLQ  
 DFDKSNPIVLRMMNDQLMFLERAFIDPLGLPDRPFYRHVIYAPSSHNKYAGESFPGIYDALFDIESKVDPSKA  
 WGEVKRQIYVAAFTVQAAAETLSEVA

## Related Products

FOLH1(PSMA)	
<a href="#">Cynomolgus_FOLH1(PSMA) CHO-K1 Cell Line</a>	<a href="#">H_FOLH1(PSMA) CHO-K1 Cell Line</a>
<a href="#">H_FOLH1(PSMA) HEK-293 Cell Line</a>	<a href="#">H_FOLH1(PSMA) RM-1 Cell Line</a>
<a href="#">Anti-FOLH1(PSMA) hIgG1 Antibody(Rosopatamab)</a>	<a href="#">Anti-FOLH1(PSMA) hIgG1 Reference Antibody (Rosobio)</a>
<a href="#">Cynomolgus FOLH1(PSMA) Protein; His Tag</a>	
ADC Related Product	
<a href="#">Anti-DXD Mouse IgG1 Antibody (23E21C5)</a>	<a href="#">Anti-DXD Mouse IgG1 Antibody (4A5A12)</a>
<a href="#">Anti-Dxd Mouse IgG2a Antibody (17D6A4)</a>	<a href="#">Anti-Eribulin Mouse IgG2a Antibody (10F8G4)</a>
<a href="#">Anti-MMAE Mouse IgG1 Antibody (11C10E3)</a>	<a href="#">Anti-MMAE Mouse IgG2a Antibody (17A1K11)</a>
<a href="#">Anti-MMAE Mouse IgG2a Antibody (8F6A3)</a>	<a href="#">Anti-SN38 Mouse IgG1 Antibody(59H11C7)</a>
<a href="#">Mouse anti Human IgG1-DXD(Dar8)</a>	<a href="#">Mouse anti Human IgG1-MMAE(Dar4)</a>
<a href="#">Human IgG1 Isotype-DXD (Dar8)</a>	<a href="#">Human IgG1 Isotype-Eribulin (Dar4)</a>
<a href="#">Human IgG1 Isotype-MMAE (Dar4)</a>	
<a href="#">Recombinant DT3C Protein</a>	
KLK2 KLK3	
<a href="#">Membrane bound H_KLK2(AA19-261) CHO-K1 Cell Line</a>	<a href="#">Membrane bound H_KLK2(AA19-261) HEK-293 Cell Line</a>
<a href="#">Membrane bound H_KLK2(AA19-261) LnCap Cell Line</a>	<a href="#">Membrane bound H_KLK2(AA25-261) CHO-K1 Cell Line</a>
<a href="#">Membrane bound H_KLK2(AA25-261) CT26 Cell Line</a>	<a href="#">Membrane bound H_KLK2(AA25-261) HEK-293 Cell Line</a>
<a href="#">Membrane bound H_KLK2(AA25-261) MC38 Cell Line (Low Expression)</a>	
<a href="#">Anti-KLK2 hIgG1 Antibody(Hu11B6)</a>	
<a href="#">Biotinylated Human KLK2 Protein; His-Avi Tag</a>	
ACP3	
<a href="#">Flag-H_ACP3 HCT116 Cell Line</a>	<a href="#">Flag-H_ACP3 HT-1080 Cell Line</a>
<a href="#">H_ACP3 HCT116 Cell Line</a>	<a href="#">H_ACP3 HT-1080 Cell Line</a>
<a href="#">H_ACP3 PC-3 Cell Line</a>	

## License Agreement:

**By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:**

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.

- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.

Genomeditech