

# Anti-MICA/MICB hIgG1 Antibody(36 NF G236A)

## Product information

GM-48843AB-10	10 µg
GM-48843AB-100	100 µg
GM-48843AB-1000	1 mg

## Antibody Information


Species Reactivity	Human
Clone	MICA.36-IgG1-NF-G236A
Source/Isotype	Monoclonal Human IgG1 /k
Application	Flow Cytometry
Specificity	Detects MICA and MICB.
Gene	MICA and MICB
Other Names	MIC-A,PERB11.1; PERB11.2
Gene ID	MICA: 100507436(human) MICB: 4277(human)
Background	<p>MHC class I polypeptide-related sequence A (MICA) is a highly polymorphic cell surface glycoprotein encoded by the MICA gene located within MHC locus. MICA is related to MHC class I and it has similar domain structure; however, it is not associated with <math>\beta</math>2-microglobulin nor binds peptides as conventional MHC class I molecules do MICA rather functions as a stress-induced ligand (as a danger signal) for integral membrane protein receptor NKG2D ("natural-killer group 2, member D").</p> <p>MHC class I polypeptide-related sequence B (MICB) is a protein that is encoded by the MICB gene located within MHC locus. MICB is related to MHC class I and has similar domain structure, which is made up of external <math>\alpha</math>1<math>\alpha</math>2<math>\alpha</math>3 domain, transmembrane segment and C-terminal cytoplasmic tail. MICB is a stress-induced ligand for NKG2D receptor. The heat shock stress pathway is involved in the regulation of MICB expression as transcription of MICB is regulated by promoter heat shock element</p>
Storage	Store at 2-8°C short term (1-2 weeks).Store at $\leq$ -20°C long term.Avoid repeated freeze-thaw.
Formulation	Phosphate-buffered solution, pH 7.2.
Endotoxin	< 1 EU/mg, determined by LAL gel clotting assay

Version:3.1 Revision Date:12/25/2023

## Data Examples

### Flow cytometry

The recommended usage range is 0.5-4  $\mu\text{g}$  per test. H\_MICA CHO-K1 Cell Line (Catalog # GM-C22346) was stained with Anti-MICA/MICB hIgG1 Antibody (Catalog # GM-48843AB) or isotype control antibody, followed by anti-Human IgG APC-conjugated Secondary Antibody.




SampleID	Geometric Mean : RL1-H
CHO-K1 anti-MICA/MICB+APC-2nd Ab	94.5
CHO-K1 H_MICA H_IgG+APC-2nd Ab	157
CHO-K1 H_MICA anti-MICA/MICB+APC-2nd Ab	34972

Fig 1. FACS

### Flow cytometry

The recommended usage range is 0.5-4  $\mu\text{g}$  per test. H\_MICA HEK-293 Cell Line (Catalog # GM-C24921) was stained with Anti-MICA/MICB hIgG1 Antibody (Catalog # GM-48843AB) or isotype control antibody, followed by anti-Human IgG APC-conjugated Secondary Antibody.



SampleID	Geometric Mean : RL1-H
HEK-293 anti-MICA/MICB+APC-2nd Ab	3704
HEK-293 H_MICA H_IgG+APC-2nd Ab	63.2
HEK-293 H_MICA anti-MICA/MICB+APC-2nd Ab	32586

Fig 2. FACS

Flow cytometry

The recommended usage range is 0.5-4 µg per test. H\_MICB HEK-293 Cell Line (Catalog # GM-C24872) was stained with Anti-MICA/MICB hlgG1 Antibody (Catalog # GM-48843AB) or isotype control antibody, followed by anti-Human IgG APC-conjugated Secondary Antibody.




Fig 3. FACS