

Product Sheet

Mouse_TLR7 Reporter 293 Cell Line


Catalog number: GM-C15439

Version 3.3.1.250117

TLR7 (Toll-like receptor 7) is an intracellular pattern recognition receptor that primarily detects single-stranded RNA (ssRNA), particularly viral RNA. It is expressed in immune cells like dendritic cells, macrophages, and B cells, contributing to the innate immune response and antiviral immunity by activating the immune system and promoting inflammatory responses and interferon production.

The TLR7 signaling pathway is activated when ssRNA binds to TLR7, recruiting the adaptor protein MyD88, which triggers downstream signaling. This pathway activates transcription factors such as NF- κ B and IRF7, leading to the expression of interferons and other cytokines. Thus, TLR7 is crucial for regulating immune responses and antiviral defense.

Mouse_TLR7 Reporter 293 Cell Line is a clonal stable HEK-293 cell line constructed using lentiviral technology, constitutive expression of the mouse TLR7 gene, along with signal-dependent expression of a luciferase reporter gene. When Resiquimod binds to TLR7, it activates downstream signaling pathways, leading to the expression of luciferase. The luciferase activity measurement indicates the activation level of the signaling pathway and can thus be used to evaluate the in vitro effects of drugs related to TLR7.



Specifications

| | |
|------------------------------|--|
| Quantity | 5E6 Cells per vial, 1 mL |
| Product Format | 1 vial of frozen cells |
| Shipping | Shipped on dry ice |
| Storage Conditions | Liquid nitrogen immediately upon receipt |
| Recovery Medium | EMEM(ATCC)+10% FBS+1% P.S |
| Growth medium | EMEM(ATCC)+10% FBS+1% P.S+3 µg/mL Blasticidin+1.5 µg/mL Puromycin |
| Note | Cells should be cultured using ATCC/30-2003 EMEM medium or Growth medium from Genomeditech. The serum should be Cegrogen biotech/A0500-3010 or sourced from Gibco. |
| Freezing Medium | 90% FBS+10% DMSO |
| Growth properties | Adherent |
| Growth Conditions | 37°C, 5% CO ₂ |
| Mycoplasma Testing | The cell line has been screened to confirm the absence of Mycoplasma species. |
| Safety considerations | Biosafety Level 2 |
| Note | It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use. |

Materials

| Reagent | Manufacturer/Catalogue No. |
|---|---|
| EMEM | ATCC/30-2003 |
| Fetal Bovine Serum | Cegrogen biotech/A0500-3010 |
| Pen/Strep | Thermo/15140-122 |
| Blasticidin | Genomeditech/ GM-040404 |
| Puromycin | Genomeditech/ GM-040401 |
| Resiquimod | MCE/HY-13740 |
| GMOne-Step Luciferase Reporter Gene Assay Kit | Genomeditech/ GM-040503 |

Figures




Figure 1 | Response to Resiquimod. Mouse_TLR7 Reporter 293 Cell Line (Cat. GM-C15439) at a concentration of 1.5E4 cells/well (96-well format) was stimulated with serial dilutions of Resiquimod (MCE/HY-13740) in assay buffer (EMEM(ATCC) + 1% FBS + 1% P.S) for 7 hours. The firefly luciferase activity was measured using the GMOne-Step Luciferase Reporter Gene Assay Kit (Cat. GM-040503). The maximum induction fold was approximately [25.8]. Data are shown by drug mass concentration.




Figure 2 | Response to Resiquimod. Mouse_TLR7 Reporter 293 Cell Line (Cat. GM-C15439) at a concentration of 1.5E4 cells/well (96-well format) was stimulated with serial dilutions of Resiquimod (MCE/HY-13740) in assay buffer (EMEM(ATCC) + 1% FBS + 1% P.S) for 7 hours. The firefly luciferase activity was measured using the GMOne-Step Luciferase Reporter Gene Assay Kit (Cat. GM-040503). The maximum induction fold was approximately [25.8]. Data are shown by drug molar concentration.

Cell Recovery

Recovery Medium: EMEM(ATCC)+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately $176 \times g$ for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at $176 \times g$ for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5×10^6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: EMEM(ATCC)+10% FBS+1% P.S+3 $\mu\text{g}/\text{mL}$ Blasticidin+1.5 $\mu\text{g}/\text{mL}$ Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at 37°C .

Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended

Medium Renewal: Every 2 to 3 days

Notes

- Upon initial revival, a higher number of dead cells and poor adherence are observed, which is normal. Adherence typically recovers within 2 - 3 days. After 2 - 3 passages, the proportion of adherent cells increases, and the cells begin to spread normally.
- After each passage, there may be 5 - 10% dead cells; however, as the number of passages increases, the recovery rate accelerates, the proportion of dead cells decreases, and the cell growth rate stabilizes.
- It is recommended to retain cell images after revival and during each observation to assist in assessing cell status. In case of abnormalities, promptly communicate with Genomeditech sales.

Related Products

| | |
|--|---|
| TLR7 | |
| H_TLR7 Reporter 293 Cell Line | |
| TLR9 | |
| H_TLR9 Reporter 293 Cell Line | Mouse_TLR9 Reporter 293 Cell Line |
| TLR8 | |
| H_TLR8 Reporter 293 Cell Line | H_TLR8 HEK-293 Cell Line |
| STING | |
| H_STING KO THP1 Cell Line | H_STING KO U937 Cell Line |
| STING KO Reporter THP1 Cell Line | STING Reporter HEK-293 Cell Line |
| STING Reporter THP1 Cell Line | STING Reporter U937 Cell Line |

Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.