

Product Sheet

H_GLP1R β-Arrestin Reporter CHO-K1 Cell Line

Catalog number: GM-C38752

Version 3.3.1.250417

The GLP-1 receptor (GLP-1R) is a G protein-coupled receptor encoded by the GLP1R gene on chromosome 6. It is expressed in pancreatic cells and brain neurons, playing a key role in insulin synthesis, secretion, and appetite regulation. GLP-1R activation increases intracellular cAMP levels via the adenylyl cyclase (AC) pathway, activating protein kinase A (PKA). This makes GLP-1R a critical target for diabetes drugs (GLP-1R agonists).

Upon agonist activation, GPCRs are phosphorylated by GRKs, exposing β -arrestin binding sites. β -arrestin binds to the receptor and cell membrane, where it stabilizes and facilitates receptor trafficking to clathrin-coated pits (CCPs) for endocytosis. This process not only desensitizes GPCRs but also activates β -arrestin-mediated, non-G protein signaling pathways.

 $H_{GLP1R} \beta$ -Arrestin Reporter CHO-K1 Cell Line is a clonal stable CHO cell line constructed using lentiviral technology, constitutive expression of the GLP1R receptor gene, and detected using enzyme fragment complementation (EFC) technology. When the agonist binds to the ligand, the two luciferase fragments complement each other to form an active complex. Upon addition of a luciferase substrate, the complex catalyzes the reaction to produce a detectable luminescent signal. This highly sensitive method enables real-time monitoring of receptor interactions as well as functional changes in the cellular environment, making it a powerful tool for drug screening and biological research.





Specifications

| Quantity | 5E6 Cells per vial,1 mL | | |
|-----------------------|--|--|--|
| Product Format | 1 vial of frozen cells | | |
| Shipping | Shipped on dry ice | | |
| Storage Conditions | Liquid nitrogen immediately upon receipt | | |
| Recovery Medium | F12K+10% FBS+1% P.S | | |
| Growth medium | F12K+10% FBS+1% P.S+4 µg/mL Puromycin | | |
| Note | None | | |
| Freezing Medium | 90% FBS+10% DMSO | | |
| Growth properties | Adherent | | |
| Growth Conditions | 37°C, 5% CO ₂ | | |
| Mycoplasma Testing | The cell line has been screened to confirm the absence of Mycoplasma species. | | |
| Safety considerations | Biosafety Level 2 | | |
| Note | It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use. | | |
| Materials | | | |

Materials

| OSTER/PYG0036 |
|---------------------------|
| grogen biotech/A0500-3010 |
| ermo/15140-122 |
| nomeditech/GM-040401 |
| CE/HY-P0055A |
| nomeditech/GM-51168AB |
| nomeditech/GM-84914AB |
| |



Figures



Figure 1 | Response to GLP-1(7-37). The H_GLP1R β -Arrestin Reporter CHO-K1 Cell Line (Cat. GM-C38752) at a concentration of 1E4 cells/well (96-well format) was stimulated with serial dilutions of GLP-1(7-37) (MCE/HY-P0055A) in assay buffer (F12K+1% FBS+1% P.S) for 16 hours. The firefly luciferase activity was measured using a detection kit. The maximum induction fold was approximately [35.2]. Data are shown by drug molar concentration.



Figure 2 | Response to Anti-GLP1R hIgG1 Antibody(mAb-36986). Serial dilutions of the Anti-GLP1R hIgG1 Antibody(mAb-36986) (Cat. GM-51168AB) was incubated with 1E4 cells/well of the H_GLP1R β -Arrestin Reporter CHO-K1 Cell Line (Cat. GM-C38752) in a 96-well plate for 1 hour in assay buffer (F12K+1% FBS+1% P.S). Subsequently, the GLP-1(7-37) (MCE/HY-P0055A) at a concentration of 8.7 nM was added, and the coculture proceeded for an additional 16 hours. The firefly luciferase activity was measured using a detection kit. The maximum induction fold was approximately [31.4]. Data are shown by drug mass concentration.

吉满生物科技(上海)有限公司 Genomeditech (Shanghai) Co., Ltd

上海市浦东新区康威路 299 号 1 幢东区 505-507 邮编 201315 505-507,5th Floor, East District, Building 1, No.299 Kangwei Road, Pudong New Area, Shanghai 本公司产品仅供科研用途,严禁用于人体治疗! For research use only!





| SampleID | Geometric Mean : FL11-H |
|---|-------------------------|
| CHO anti-GLP1R+APC-2nd Ab | 1311 |
| H_GLP1R β-Arrestin Reporter CHO H_IgG+APC-2nd Ab | 1190 |
| H_GLP1R β-Arrestin Reporter CHO anti-GLP1R+APC-2nd Ab | 44517 |

Figure 3 | H_GLP1R β-Arrestin Reporter CHO-K1 Cell Line (Cat. GM-C38752) was determined by flow cytometry using Anti-H_GLP1R hIgG1 Antibody (glutazumab) (Cat. GM-84914AB).

Cell Recovery

Recovery Medium: F12K+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70° C. Storage at -70° C will result in loss of viability.

- a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 3 minutes).
- b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO₂ in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at 176 x g for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

吉满生物科技(上海)有限公司 Genomeditech (Shanghai) Co., Ltd



Cell passage

Growth medium: F12K+10% FBS+1% P.S+4 µg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Remove and discard culture medium.
- b) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- c) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 3 minutes at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach.
 Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- e) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- f) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- g) Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:4 - 1:5 is recommended

Medium Renewal: Every 2 to 3 days

Notes

a) After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

| GCGR | | | | |
|--|---|--|--|--|
| H_GCGR Reporter CHO-K1 Cell Line | H_GCGR Reporter HEK-293 Cell Line | | | |
| H_GCGR Reporter HEK-293 DDX35TM Cell Line | Cynomolgus_GCGR HEK-293 Cell Line | | | |
| H_GCGR CHO-K1 Cell Line | H_GCGR HEK-293 Cell Line | | | |
| Mouse_GCGR HEK-293 Cell Line | | | | |
| Anti-H_GCGR hIgG2 Antibody(volagidemab) | | | | |
| GLP1R | | | | |
| H_GLP1R Reporter CHO-K1 Cell Line | H_GLP1R Reporter HEK-293 Cell Line | | | |
| H_GLP1R Reporter HEK-293 DDX35TM Cell Line | Cynomolgus_GLP1R HEK-293 Cell Line | | | |
| H_GLP1R CHO-K1 Cell Line | H_GLP1R HEK-293 Cell Line | | | |
| Mouse_GLP1R HEK-293 Cell Line | | | | |
| Anti-GLP1R hIgG1 Antibody(mAb-36986) | Anti-H_GLP1R hIgG1 Antibody(glutazumab) | | | |
| FGF21 | | | | |
| H_FGF21 Reporter HEK-293 Cell Line | | | | |
| CALCA(CGRP): CALCRL RAMP | | | | |
| H_CALCRL RAMP1 Reporter HEK-293 Cell Line | H_CALCRL RAMP1 Reporter HEK-293 DDX35TM Cell Line | | | |

Related Products

吉满生物科技(上海)有限公司 Genomeditech (Shanghai) Co., Ltd

上海市浦东新区康威路 299 号 1 幢东区 505-507 邮编 201315 505-507,5th Floor, East District, Building 1,No.299 Kangwei Road, Pudong New Area, Shanghai 本公司产品仅供科研用途,严禁用于人体治疗! For research use only!

Genomeditech

Genomeditech (Shanghai) Co.,Ltd. Order: +86 021-68455258/50432826/50432825 Toll-free: +86 400 627 9288 Email: service@genomeditech.com

| Cynomolgus_CALCRL RAMP1 HEK-293 Cell Line | H_CALCRL RAMP1 CHO-K1 Cell Line | | | |
|--|--|--|--|--|
| H_CALCRL RAMP1 HEK-293 Cell Line | | | | |
| Anti-CALCRL RAMP1 hIgG2 Antibody(Erenumab) | | | | |
| GIP:GIPR | | | | |
| H_GIPR Reporter CHO-K1 Cell Line | H_GIPR Reporter HEK-293 Cell Line | | | |
| H_GIPR Reporter HEK-293 DDX35TM Cell Line | Cynomolgus_GIPR HEK-293 Cell Line | | | |
| H_GIPR CHO-K1 Cell Line | H_GIPR HEK-293 Cell Line | | | |
| Mouse_GIPR HEK-293 Cell Line | | | | |
| Anti-H_GIPR hIgG1 Antibody(AMG-133) | | | | |
| ACVR2A: ACTRIIB: Active A | | | | |
| ACVR2A KO HEK-293 Cell Line | Activin A Reporter Cell Line | | | |
| BRE Reporter 293 Cell Line | H_ACVR2A Reporter Cell Line | | | |
| H_ACVR2B Reporter Cell Line | ACVR2B KO HEK-293 Cell Line | | | |
| H_ACVR2A HEK-293(ACVR2B KO) Cell Line | H_ACVR2B CHO-K1 Cell Line | | | |
| H_ACVR2B HEK-293(ACVR2A KO) Cell Line | | | | |
| Anti-ACVR2B hIgG1 Antibody(Bimagrumab) | Anti-ACVR2B hIgG1 Antibody(Fab-17G05) | | | |
| Anti-ACVR2B mIgG2a Antibody(Bimagrumab) | Anti-H_ACVR2B hIgG1 Reference Antibody(Bimbio) | | | |
| Biotinylated Human ACVR2A Protein; His-Avi Tag | Biotinylated Human ACVR2B Protein; His-Avi Tag | | | |
| Biotinylated Mouse ACVR2A Protein; His-Avi Tag | Biotinylated Mouse ACVR2B Protein; His-Avi Tag | | | |
| Human Activin A Protein; His Tag | Human Activin B Protein; His Tag | | | |
| Human ACVR2A Protein; hFc Tag | Human ACVR2A Protein; His Tag | | | |
| Human ACVR2B Protein; hFc Tag | Human ACVR2B Protein; His Tag | | | |
| Human latent GDF-8 Protein; His Tag | Mouse ACVR2A Protein; His Tag | | | |
| Mouse ACVR2B Protein; His Tag | | | | |
| AMY: CALCR RAMP | | | | |
| H_CALCR RAMP3(AMY3) Reporter CHO-K1 Cell Line | H_CALCR Reporter CHO-K1 Cell Line | | | |

Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.

吉满生物科技(上海)有限公司 Genomeditech (Shanghai) Co., Ltd

上海市浦东新区康威路 299 号 1 幢东区 505-507 邮编 201315 505-507,5th Floor, East District, Building 1, No.299 Kangwei Road, Pudong New Area, Shanghai 本公司产品仅供科研用途,严禁用于人体治疗! For research use only!