

Product Sheet

H_RXFP1 Reporter 293 Cell Line


Catalog number: GM-C42543

Version 3.3.1.260330

RXFP1 (Relaxin Family Peptide Receptor 1, alias LGR7) belongs to the Relaxin/Insulin-like Peptide Receptor Family (RXFPs) and is a member of the Class A subfamily of the G protein-coupled receptor (GPCR) superfamily. It recognizes ligands such as relaxin through its extracellular leucine-rich repeat (LRR) domain and mediates signaling via G protein coupling through its seven-transmembrane domain. Its gene is located on human chromosome 4q32.1, encoding a protein of 757 amino acids. RXFP1 is widely expressed in uterine smooth muscle, cardiomyocytes, vascular endothelium, kidneys, and macrophages, where it regulates reproduction (pregnancy maintenance, cervical softening), cardiovascular functions (anti-fibrosis, vasodilation), and tissue repair (anti-inflammation, wound healing).

H_RXFP1 Reporter 293 Cell Line is a clonal stable 293 cell line constructed using lentiviral technology, constitutive expression of the RXFP1 gene, along with signal-dependent expression of a luciferase reporter gene. When Relaxin-2, ML-290, or AZD5462 binds to the receptor, the downstream signaling pathway is activated, thereby inducing luciferase expression. The luciferase activity measurement indicates the activation level of the signaling pathway and can thus be used to evaluate the in vitro effects of drugs related to RXFP1.

Relaxin-2/ML-290/AZD5462



Specifications

Quantity	5E6 Cells per vial, 1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Recovery Medium	DMEM+10% FBS+1% P.S
Growth medium	DMEM+10% FBS+1% P.S+4 µg/mL Blasticidin+0.75 µg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO ₂
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Blasticidin	Genomeditech/ GM-040404
Puromycin	Genomeditech/ GM-040401
Human Relaxin-2 Recombinant Protein	Thermofisher/130-15-25UG
ML-290	MCE/HY-112606
AZD5462	Bidepharm/BD01590549
GMOne-Step 2.0 Luciferase Reporter Gene Assay Kit	Genomeditech/ GM-040513

Figures




Figure 1 | Response to Human Relaxin-2 Recombinant Protein. The H_RXFP1 Reporter 293 Cell Line (Cat. GM-C42543) at a concentration of 1.5E4 cells/well (96-well format) was stimulated with serial dilutions of Human Relaxin-2 Recombinant Protein (Thermofisher/130-15-25UG) in assay buffer (DMEM+1% FBS+1% P.S) for 16 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [68.5]. Data are shown by drug mass concentration.




Figure 2 | Response to Human Relaxin-2 Recombinant Protein. The H_RXFP1 Reporter 293 Cell Line (Cat. GM-C42543) at a concentration of 1.5E4 cells/well (96-well format) was stimulated with serial dilutions of Human Relaxin-2 Recombinant Protein (Thermofisher/130-15-25UG) in assay buffer (DMEM+1% FBS+1% P.S) for 16 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [68.5]. Data are shown by drug molar concentration.




Figure 3 | Response to ML-290. The H_RXFP1 Reporter 293 Cell Line (Cat. GM-C42543) at a concentration of 1.5E4 cells/well (96-well format) was stimulated with serial dilutions of ML-290 (MCE/HY-112606) in assay buffer (DMEM+1% FBS+1% P.S) for 16 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [52.8]. Data are shown by drug mass concentration.




Figure 4 | Response to AZD5462. The H_RXFP1 Reporter 293 Cell Line (Cat. GM-C42543) at a concentration of 1.5E4 cells/well (96-well format) was stimulated with serial dilutions of AZD5462 (Bidepharm/BD01590549) in assay buffer (DMEM+1% FBS+1% P.S) for 16 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [51.9]. Data are shown by drug mass concentration.




Figure 5 | The mRNA expression levels of H_RXFP1 in the H_RXFP1 Reporter 293 Cell Line (Cat. GM-C42543) were determined by RT-qPCR.

Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately $176 \times g$ for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at $176 \times g$ for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5×10^6 cells/mL.
- Aliquot 1 mL into each vial.

- d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: DMEM+10% FBS+1% P.S+4 $\mu\text{g}/\text{mL}$ Blasticidin+0.75 $\mu\text{g}/\text{mL}$ Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Subculturing is necessary when the cell density reaches 80%. It is recommended to perform subculturing at a ratio of 1:3 to 1:4 every 2-3 days. Ensure that the density does not exceed 80%, as overcrowding can lead to reduced viability due to compression.
- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at 37°C .

Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended

Medium Renewal: Every 2 to 3 days

Notes

- Upon initial thawing, a higher number of dead cells is observed, which is a normal phenomenon. Significant improvement is seen after adaptation. Once the cells reach a stable state, the number of dead cells decreases after subculturing and the cell growth rate becomes stable.
- Ensure that the cell density does not exceed 80%, as overcrowding may lead to reduced viability due to compression.

Related Products

APJ	
H_APJ Reporter HEK-293 Cell Line	
NPR1	
H_NPR1 Reporter Cell Line	Cynomolgus_NPR1 CHO-K1 Cell Line
Flag-Mouse_NPR1 CHO-K1 Cell Line	H_NPR1 CHO-K1 Cell Line
H_NPR1 HEK-293 Cell Line	Mouse_NPR1 CHO-K1 Cell Line
Rat_NPR1 CHO-K1 Cell Line	
Anti-NPR1 hIgG1 Antibody(XX-16)	Anti-NPR1 hIgG1 Reference Antibody (XX-16)

Anti-NPR1 hIgG4 Antibody(REGN-5381)	Anti-NPR1 hIgG4 Reference Antibody (REGN-5381)
NPR2	
H_NPR2 Reporter 293 Cell Line	

License Agreement:

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.

Genomeditech