

Product Sheet



H_GUCY2C HCT116 Cell Line

Catalog number: GM-C26271

Version 4.0.26.924

Specifications


Description	H_GUCY2C HCT116 Cell Line is a clonal stable HCT116 cell line that constitutively expresses the human GUCY2C gene, constructed using lentiviral technology.
Quantity	5E6 Cells per vial, 1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Target	Human GUCY2C
Gene ID/Uniprot ID	P25092
Host Cell	HCT116
Recovery Medium	McCoy's 5A+10% FBS+1% P.S
Growth medium	McCoy's 5A+10% FBS+1% P.S+0.5 µg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO ₂
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
McCoy's 5A	VivaCell/C3020-0500
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/ GM-040401
Human Guanylyl Cyclase C/GUCY2C Antibody (2543C)	R&D Systems/MAB2157
Countess 3 Automated Cell Counter	ThermoFisher/A49891
NovoCyt Flow Cytometer	Agilent/NovoCyt 2060R

Data

Expression



	SampleID	Geometric Mean : R675-H
	HCT116 anti-GUCY2C+APC-2nd Ab	823
	HCT116 H_GUCY2C Rabbit_IgG+APC-2nd	854
	HCT116 H_GUCY2C anti-GUCY2C+APC-2nd Ab	14501

H_GUCY2C HCT116 Cell Line (Cat. GM-C26271) was determined by flow cytometry using Human Guanylyl Cyclase C/GUCY2C Antibody (2543C)(R&D Systems/MAB2157).

Cell Recovery

Recovery Medium: McCoy's 5A+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO₂ in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at 176 x g for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: McCoy's 5A+10% FBS+1% P.S+0.5 µg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Remove and discard culture medium.
- b) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- c) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 1 to 2 minutes at 37°C).
- d) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- e) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- f) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- g) Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended

Medium Renewal: Every 1 to 2 days

Notes

- a) After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

Sequence

GUCY2C [P25092](#)

MKTLLLDLALWSLLFQPGWLSFSSQVSQNCHNGSYEISVLMMGNSAFAEPLKNLEDAVNEGLEIVRGRLQNAGL
NVTVNATFMYS DGLIHNSGDCRSSTCEGLDLLRKISNAQRMGCVLIGPSCYSTYTFQMYLDTELSYPMISAGSFGLS
CDYKETLTRLMS PARKLMYFLVNFWKTNLDPFKTYSWSTSYVYKNGTETEDCFWYLNALASVSYSFHELGFK
VVLQRQDKEFQDILMDHNRKSNVIMCGGPEFLYKLGDRABAEDIVILVDFLNDQYFEDNVTAPDYMKNVLVL
TLSPGNSLLNSSFSRNLSPTKRDFALAYLNGILLFGHMLKIFLENGENITTPKFAHAFRNLTFEGYDGPVTLDDWG
DVDSTMVLLYTSVDTKKYKVLITYDTHVNKTYPVDMSPFTWKNKLPNDITGRGPQILMIAVFTLTGAVVLLL
LVALLMLRKYRKDYELRQKKWSHIPPENIFPLETNETNHVSLKIDDDKRRDTIQRLRQCKYDKKRVLKDLKHN
DGNFTEKQKIELNKLQIDYYNLTKFYGTVKLDTMIFGVIEYCERGLREVNDTISYPDGTFMDWEFKISVLYDI
AKGMSYLHSSKTEVHGRLKSTNCVVD SRMVVKITDFGCNSILPPKKDLWTAPEHLRQANISQKGDVYSYGIIAQE
IILRKETFYTLSCRDRNEKIFRVENSNGMKPFRPDLFLETAEEKELEVYLLVKNCWEEDPEKRPDFKKIETTLAKIF
GLFHDQKNESYMDTLIRRLQLYSRNLEHLVEERTQLYKAERDRADRLNFMLLPRLVVKSLKEKGFVEPELYEEV
TIYFSDIVGFTTICKYSTPMEVVDMLNDIYKSFDHIVDHHDVYKVETIGDAYMVASGLPKRNGNRHAIDIAKMAL
EILSFMGTFELEHLPGLPIWIRIGVHSGPCAAGVVGIKMPRYCLFGDTVNTASRMESTGLPLRIHVSGSTIAILKRTE
CQFLYEVRGETYLGKRGNETTYWLTGMKDQKFNLPTPTVENQQRLQAEFSDMIANS LQKRQAAGIRSQKPRR
VASYKKGTLEYLQLNTTDDKESTYF

Related Products

CLDN18	
Cynomolgus_CLDN18.2-eGFP CHO-K1 Cell Line	H_CLDN18(isoform2)-eGFP 293 Cell Line
H_CLDN18.1-eGFP HEK-293 Cell Line	H_CLDN18.2 MC38 Cell Line
H_CLDN18.2 MKN45 Cell Line(High Expression)	H_CLDN18.2 MKN45 Cell Line(Low Expression)
H_CLDN18.2 MKN45 Cell Line(Medium Expression)	H_CLDN18.2 PANC-1 Cell Line
H_CLDN18.2(isoform2) CHO-K1 Cell Line	H_CLDN18.2-eGFP CT-26 Cell Line
Mouse_CLDN18.2-eGFP CHO-K1 Cell Line	Rat_CLDN18.2-eGFP CHO-K1 Cell Line
Rhesus_CLDN18.2-eGFP CHO-K1 Cell Line	
Anti-CLDN18.2 hIgG1 Reference Antibody (IMAB362)	Anti-CLDN18.2 hIgG1 Antibody(LM-102)
Anti-CLDN18.2 hIgG1 Antibody(Zolbetuximab)	
TROP2(TACSTD2)	
Cynomolgus_Trop2 CHO-K1 Cell Line	Cynomolgus_TROP2 HEK-293 Cell Line
H_TROP2 CHO-K1 Cell Line	H_TROP2 CT26 Cell Line
H_TROP2 HEK-293 Cell Line	H_TROP2 LLC1 Cell Line
Anti-H_TROP2 hIgG1 Antibody(Datopotamab)	Anti-TROP2 hIgG1 Antibody(Hu2G10-5)
Anti-Trop2 hIgG1 Reference Antibody (Sacbio)	Anti-Trop2 hIgG1 Reference Antibody(Datbio)

Anti-Trop2-DXD ADC(Dar4)[Datopotamab deruxtecán,Dato-DXD]	Anti-Trop2-SN38 ADC(Dar8)[Sacituzumab govitecan]
Biotinylated Cynomolgus TROP2 Protein; His-Avi Tag	Biotinylated Human TROP2 Protein; His-Avi Tag
Human TROP2 Protein; hFc Tag	Human TROP2 Protein; His Tag
GUCY2C(GC-C)	
Cynomolgus_GUCY2C HEK-293 Cell Line	H_GUCY2C CHO-K1 Cell Line
H_GUCY2C HEK-293 Cell Line	
Anti-H_GUCY2C hIgG1 Antibody(Indusatumab)	
Cynomolgus GUCY2C Protein; His Tag	Human GUCY2C Protein; His Tag
FGFR2	
H_FGFR2b Reporter 293 Cell Line	Cynomolgus_FGFR2b HEK-293 Cell Line
H_FGFR2b CHO-K1 Cell Line	H_FGFR2b HEK-293 Cell Line
Mouse_FGFR2b HEK-293 Cell Line	
Anti-FGFR2 hIgG1 Antibody(hFR2-14_H12-L1)	Anti-FGFR2 hIgG1 Reference Antibody (Bemabio)
Cynomolgus FGFR2(IIIb) D1-D3 Protein; His Tag	Human FGFR2(IIIb) D1-D3 Protein; hFc Tag
Human FGFR2(IIIb) D2-D3 Protein; hFc Tag	Human FGFR2(IIIb) D2-D3 Protein; His Tag
Human FGFR2(IIIc) D1-D3 Protein; His Tag	Mouse FGFR2(IIIb) D2-D3 Protein; His Tag
CD44	
H_CD44s CHO-K1 Cell Line	H_CD44v6 CHO-K1 Cell Line
H_CD44v6 CHO-K1 Cell Line (Medium Expression)	H_CD44v6 HEK-293 Cell Line
Anti-CD44v6 hIgG1 Antibody(bivatuzumab)	
ADC Related Product	
Anti-DXD Mouse IgG1 Antibody (23E21C5)	Anti-DXD Mouse IgG1 Antibody (4A5A12)
Anti-Dxd Mouse IgG2a Antibody (17D6A4)	Anti-Eribulin Mouse IgG2a Antibody (10F8G4)
Anti-MMAE Mouse IgG1 Antibody (11C10E3)	Anti-MMAE Mouse IgG2a Antibody (17A1K11)
Anti-MMAE Mouse IgG2a Antibody (8F6A3)	Anti-SN38 Mouse IgG1 Antibody(59H11C7)
Mouse anti Human IgG1-DXD(Dar8)	Mouse anti Human IgG1-MMAE(Dar4)
Human IgG1 Isotype-DXD (Dar8)	Human IgG1 Isotype-Eribulin (Dar4)
Human IgG1 Isotype-MMAE (Dar4)	
Recombinant DT3C Protein	

License Agreement

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.